

Stroke Aphasia Depression Questionnaire (SADQ-10) – Community Version

Reproduced with permission: Sutcliffe, L. & Lincoln, N. (1998). The assessment of depression in aphasic stroke patients: The development of the Stroke Aphasic Depression Questionnaire. *Clinical Rehabilitation*, 12, 506-513.

The community SADQ-IO provides community based health care workers with a structured and reliable approach to assessing the likelihood that an individual with a speech or language impairment is suffering depressed mood.

Please indicate how often in the last week the patient has shown the following behaviours:

Does he/she have weeping spells?	Sometimes	Rarely	Never
Does he/she have restless disturbed nights?	Sometimes	Rarely	Never
Does he/she avoid eye contact when you talk to him/her?	Sometimes	Rarely	Never
Does he/she burst into tears?	Sometimes	Rarely	Never
Does he/she complain of aches and pains?	Sometimes	Rarely	Never
Does he/she get angry?	Sometimes	Rarely	Never
Does he/she refuse to participate in social activities?	Sometimes	Rarely	Never
Is he/she restless and fidgety?	Sometimes	Rarely	Never
Does he/she sit without doing anything?	Sometimes	Rarely	Never
Does he/she keep him/herself occupied during the day?	Sometimes	Rarely	Never